

Noncompact homogeneous Einstein manifolds attached to graded Lie algebras

Hiroshi Tamaru

Department of Mathematics, Hiroshima University,
1-3-1 Kagamiyama, Higashi-Hiroshima 739-8526, JAPAN
(e-mail: tamaru@math.sci.hiroshima-u.ac.jp)

Abstract

In this paper, we study the nilradicals of parabolic subalgebras of semisimple Lie algebras and the natural one-dimensional solvable extensions of them. We investigate the structures, curvatures and Einstein conditions of the associated nilmanifolds and solvmanifolds. We show that our solvmanifold is Einstein if the nilradical is of two-step. New examples of Einstein solvmanifolds with three-step and four-step nilradicals are also given.

0 Introduction

A Riemannian manifold is called a *solvmanifold* if it admits a transitive solvable group of isometries. Up to now, all known examples of noncompact homogeneous Einstein manifolds are solvmanifolds. The study of Einstein solvmanifolds is currently very active. In particular, Heber ([11]) deeply studied Einstein solvmanifolds and obtained many essential results. One of his theorems states that the moduli space of Einstein solvmanifolds may have large dimension. Despite of that, we do not know many explicit examples of Einstein solvmanifolds; in particular, Einstein solvmanifolds whose nilradicals have the nilpotency greater than two. The purpose of this paper is to introduce and study a class of solvmanifolds, and provide new examples of Einstein spaces. Our class contains Einstein solvmanifolds whose nilradicals are of two, three and four-step. We need to know explicit examples for the further study of Einstein solvmanifolds.

Here we briefly recall some known examples of noncompact homogeneous Einstein manifolds. One strategy for finding Einstein solvmanifolds is to consider solvmanifolds

⁰Mathematics Subject Classification (2000): 53C30, 22E25, 53C25, 53C35,

with good geometric structures. Typical examples are symmetric spaces of noncompact type. Other examples are noncompact homogeneous Kähler-Einstein manifolds, which are modeled by solvable normal j -algebras (see [20]), and quaternionic Kähler solvmanifolds (see [2], [7]). The other strategy is to consider solvable extensions of good nilmanifolds. Typical examples are Damek-Ricci spaces, which are the one-dimensional solvable extensions of generalized Heisenberg groups ([5], [8], see also [3]). Other examples are given by Mori ([18]); he started from complex graded Lie algebras of second kind and considered the solvable extensions of the associated nilmanifolds. In fact, our construction is a generalization of his study. Other strategy for finding Einstein solvmanifolds can be found in recent articles, for example, see [15], [16] and [10]. The list of known examples of Einstein solvmanifolds can be found in [17].

In this paper we employ the second strategy. We take the nilradicals of parabolic subalgebras of semisimple Lie algebras. These nilpotent Lie algebras admit the natural inner products, and can be investigated in term of graded Lie algebras. In Section 2, we mention some properties of graded Lie algebras and define our nilmanifolds. The structures and curvatures of our nilmanifolds will be studied in Sections 3 and 4. In Section 5, we study curvature property of one-dimensional solvable extensions of our nilmanifolds. We find that there is a natural solvable extension, that is, only one candidate to be Einstein. Curvatures of the natural solvable extension is studied in Section 6. We see in Section 7 that our solvmanifold is Einstein if the nilradical is of two-step. This provides many new examples of Einstein solvmanifolds and some of them have "unusual" eigenvalue types. Other new examples of Einstein solvmanifolds with three-step and four-step nilradicals will be given in Section 8. We observe in Section 9 that, if $\mathfrak{g}$ is small, then every attached solvmanifolds are Einstein.

Note that, if a parabolic subalgebra is minimal, then its nilradical coincides with the nilpotent part of the Iwasawa decomposition. In this case the constructed solvmanifold coincides with the rank one reduction of a symmetric space of noncompact type. Therefore our class of solvmanifolds essentially contains noncompact symmetric spaces. The structures of the rank one reductions of noncompact symmetric spaces will be mentioned in Section 10.

The author would like to thank Professor Soji Kaneyuki, Professor Hiroshi Asano and the referee. Theorem 4.3 was pointed out by Professor Asano and the referee. Theorem 5.3 was suggested by the referee. Their advices are valuable to refine the first version of this manuscript. This work was partially supported by Grant-in-Aid for Young Scientists (B) 14740049 and 17740039, The Ministry of Education, Culture, Sports, Science and Technology, Japan.

1 Preliminaries

In this section we recall some notations and basic properties of Einstein solvmanifolds. We refer to [24], [11].

Let $\mathfrak{s}$ be a solvable Lie algebra and $\langle, \rangle$ be a positive definite inner product on $\mathfrak{s}$. Throughout this paper, by a metric Lie algebra $(\mathfrak{s}, \langle, \rangle)$ we also mean the corresponding solvmanifold, that is, the simply-connected Lie group S equipped with the induced left-invariant Riemannian metric. Put $\mathfrak{n} := [\mathfrak{s}, \mathfrak{s}]$ and $\mathfrak{a} := \mathfrak{n}^\perp$.

Definition 1.1 A solvmanifold $(\mathfrak{s}, \langle, \rangle)$ is called *standard* if $\mathfrak{a}$ is abelian. A standard solvmanifold $(\mathfrak{s}, \langle, \rangle)$ is said to be of *Iwasawa-type* if (i) ad_A is symmetric for every $A \in \mathfrak{a}$, and (ii) there exists $A_0 \in \mathfrak{a}$ such that every eigenvalue of ad_{A_0} on $\mathfrak{n}$ is a positive real number.

To calculate the Ricci curvatures of our solvmanifolds, we need the following particular vectors.

Definition 1.2 For a solvmanifold $(\mathfrak{s}, \langle, \rangle)$, we call $H_0 \in \mathfrak{a}$ the *mean curvature vector* if $\langle H_0, A \rangle = \text{tr}(\text{ad}_A)$ for every $A \in \mathfrak{a}$.

Note that H_0 is the mean curvature vector for the embedding of N into S , where N denotes the Lie subgroup of S with Lie algebra $\mathfrak{n}$.

Proposition 1.3 ([24]) *Let $(\mathfrak{s}, \langle, \rangle)$ be a solvmanifold of Iwasawa type and H_0 be the mean curvature vector. Denote by Ric and $\text{Ric}^\mathfrak{n}$ the Ricci curvatures of $\mathfrak{s}$ and $\mathfrak{n}$, respectively. Then,*

- (1) $\text{Ric}(A, A') = -\text{tr}(\text{ad}_A) \circ (\text{ad}_{A'})$ for $A, A' \in \mathfrak{a}$,
- (2) $\text{Ric}(A, X) = 0$,
- (3) $\text{Ric}(X) = \text{Ric}^\mathfrak{n}(X) - (\text{tr}(\text{ad}_{H_0})/|H_0|^2) \cdot \text{ad}_{H_0}X$ for $X \in \mathfrak{n}$.

Proposition 1.3 leads that, if $(\mathfrak{s}, \langle, \rangle)$ is Einstein then so is $(\mathbb{R}H_0 + \mathfrak{n}, \langle, \rangle)$. For a solvmanifold $(\mathfrak{s}, \langle, \rangle)$, we call $\dim \mathfrak{a}$ the *algebraic rank* and $(\mathbb{R}H_0 + \mathfrak{n}, \langle, \rangle)$ the *rank one reduction*. Note that H_0 is also the mean curvature vector of the solvmanifold $(\mathbb{R}H_0 + \mathfrak{n}, \langle, \rangle)$.

Proposition 1.4 ([1]) *The Ricci curvature of a nilmanifold $(\mathfrak{n}, \langle, \rangle)$ is given by*

$$\text{Ric}^\mathfrak{n} = (1/4) \sum \text{ad}_{E_i} \circ (\text{ad}_{E_i})^* - (1/2) \sum (\text{ad}_{E_i})^* \circ \text{ad}_{E_i},$$

where $\{E_i\}$ is an orthonormal basis of $\mathfrak{n}$ and $(\text{ad}_{E_i})^*$ denotes the dual of the adjoint operator defined by $\langle (\text{ad}_{E_i})^*X, Y \rangle = \langle X, \text{ad}_{E_i}Y \rangle$.

Propositions 1.3 and 1.4 state that the Ricci curvatures of solvmanifolds of Iwasawa-type can be calculated in terms of the mean curvature vectors, adjoint operators and their dual operators.

Proposition 1.5 ([11]) *For a standard Einstein solvmanifold, there exists $\lambda > 0$ such that the real part of every eigenvalue of $\text{ad}_{\lambda H_0}$ is a positive integer.*

Let $\mu_1 < \mu_2 < \dots < \mu_m$ be the real parts of eigenvalues of $\text{ad}_{\lambda H_0}$ without common divisor, and $d_1, d_2, \dots, d_m$ the corresponding multiplicities. We call $(\mu_1 < \dots < \mu_m; d_1, \dots, d_m)$ the *eigenvalue type* of an Einstein solvmanifold. For example, the eigenvalue types of Damek-Ricci spaces are $(1 < 2; d_1, d_2)$.

2 Nilmanifolds attached to graded Lie algebras

In this section we briefly recall some properties of graded Lie algebras and define the nilmanifolds associated to them. We refer to [13], [22] for graded Lie algebras.

Let $\mathfrak{g}$ be a semisimple Lie algebra. A decomposition of $\mathfrak{g}$ into subspaces,

$$\mathfrak{g} = \sum_{k \in \mathbb{Z}} \mathfrak{g}_k$$

is called a *gradation* if $[\mathfrak{g}_k, \mathfrak{g}_l] \subset \mathfrak{g}_{k+l}$ holds for every k and l . A Lie algebra with gradation is called a *graded Lie algebra*. By definition, $\mathfrak{n} := \sum_{k > 0} \mathfrak{g}_k$ is a nilpotent subalgebra.

In the theory of graded Lie algebras, it is fundamental that there exist the characteristic element $Z \in \mathfrak{g}$ and a grade-reversing Cartan involution σ . An element Z is called the *characteristic element* if $\mathfrak{g}_k = \{X \in \mathfrak{g} \mid [Z, X] = k \cdot X\}$ for every k . A Cartan involution σ is said to be *grade-reversing* if $\sigma(\mathfrak{g}_k) = \mathfrak{g}_{-k}$ for every k . Note that the characteristic element is unique. It is easy to show that σ is grade-reversing if and only if $\sigma(Z) = -Z$. A Cartan involution σ defines the natural inner product B_σ on $\mathfrak{g}$ by $B_\sigma(X, Y) := -B(X, \sigma(Y))$, where B denotes the Killing form of $\mathfrak{g}$. Now we can define our nilmanifolds:

Definition 2.1 Let $\mathfrak{g} = \sum \mathfrak{g}_k$ be a graded Lie algebra. The nilpotent Lie algebra $\mathfrak{n} := \sum_{k > 0} \mathfrak{g}_k$ endowed with the inner product B_σ is called the *nilmanifold attached to the graded Lie algebra*, or shortly, *nilmanifold attached to GLA*.

Note that $\sum_{k \geq 0} \mathfrak{g}_k$ is a parabolic subalgebra of $\mathfrak{g}$ and $\mathfrak{n}$ is the nilradical of it. Conversely, it is known that every parabolic subalgebra of $\mathfrak{g}$ can be obtained in this way. Therefore, we are considering the nilradicals of all parabolic subalgebras of all semisimple Lie algebras.

We also note that our nilpotent Lie algebras $\mathfrak{n}$ are related to R -spaces. Let $\mathfrak{g} = \sum \mathfrak{g}_k$ be a graded Lie algebra, G the centerless Lie group with Lie algebra $\mathfrak{g}$, and P the subgroup of G with Lie algebra $\sum_{k \leq 0} \mathfrak{g}_k$. The quotient space G/P is an R -space (it is sometimes called a real flag manifold). By definition, $\mathfrak{n}$ can be naturally identified with the tangent spaces of G/P . Therefore, there is a correspondence between the class of our nilmanifolds and the class of R -spaces (but this is not one-to-one).

A graded Lie algebra $\mathfrak{g} = \sum \mathfrak{g}_k$ is said to be of ν -th kind if $\mathfrak{g}_\nu \neq 0$ and $\mathfrak{g}_k = 0$ for every $k > \nu$. The existence of a grade-reversing Cartan involution implies that $\dim \mathfrak{g}_k = \dim \mathfrak{g}_{-k}$, therefore, a ν -th kind gradation satisfies $\mathfrak{g}_k = 0$ for every $|k| > \nu$. Note that, for a graded Lie algebra of ν -th kind, the nilpotent Lie algebra $\mathfrak{n} = \sum_{k > 0} \mathfrak{g}_k$ is not of ν -step in general, but at most ν -step nilpotent. We will explain this later.

To state the classification of gradations, we now start from a semisimple symmetric pair of noncompact type $(\mathfrak{g}, \mathfrak{k})$ with Cartan involution σ . Take the Cartan decomposition $\mathfrak{g} = \mathfrak{k} + \mathfrak{p}$ with respect to σ and a maximal abelian subspace $\mathfrak{a}$ in $\mathfrak{p}$. Let Δ be the root system of $\mathfrak{g}$ with respect to $\mathfrak{a}$. Recall that we say $\alpha \in \mathfrak{a}^* - \{0\}$ is a *root* if

$$0 \neq \mathfrak{g}_\alpha := \{X \in \mathfrak{g} \mid [H, X] = \alpha(H) \cdot X \text{ for every } H \in \mathfrak{a}\}.$$

Note that $\mathfrak{g}$ can be decomposed into the direct sum of root spaces: $\mathfrak{g} = \mathfrak{g}_0 + \sum_{\alpha \in \Delta} \mathfrak{g}_\alpha$. Let $\Pi := \{\alpha_1, \dots, \alpha_r\}$ be a set of simple roots of Δ and define

$$\begin{aligned} C &:= \{H \in \mathfrak{a} \mid \alpha_i(H) \in \mathbb{Z}_{\geq 0} \text{ for every } \alpha_i \in \Pi\} \\ &= \{c_1 H^1 + \dots + c_r H^r \mid c_1, \dots, c_r \in \mathbb{Z}_{\geq 0}\}, \end{aligned}$$

where $\{H^1, \dots, H^r\}$ is the dual basis of Π , that is, $\alpha_i(H^j) = \delta_{ij}$. Then, every element $Z \in C$ gives a gradation, by putting

$$\mathfrak{g}_k := \sum_{\alpha(Z)=k} \mathfrak{g}_\alpha.$$

On the other hand, every gradation of $\mathfrak{g}$ can be obtained in this way up to conjugation (see [14], [22]). We summarize this as follows.

Theorem 2.2 ([14], [22]) *There is a bijective correspondence between the set of the isomorphism classes of gradations on $\mathfrak{g}$ and the set of $\text{Aut}(DD)$ -orbits of C , where $\text{Aut}(DD)$ denotes the automorphism group of the Dynkin diagram of Δ .*

It is easy to see that the gradation given by $Z \in C$ is of ν -th kind if and only if $\nu = \tilde{\alpha}(Z)$, where $\tilde{\alpha}$ denotes the highest root. Obviously $\mathfrak{n} = \sum_{k > 0} \mathfrak{g}_k$ is at most ν -step nilpotent. We will mention here the condition for $\mathfrak{n}$ to be ν -step nilpotent.

A graded Lie algebra $\mathfrak{g} = \sum \mathfrak{g}_k$ is said to be of *type* α_0 if $\mathfrak{g}_1$ generates $\sum_{k > 0} \mathfrak{g}_k$. This notation was introduced by Kaneyuki and Asano ([14]), and they showed that a graded Lie algebra $\mathfrak{g} = \sum \mathfrak{g}_k$ is of type α_0 if and only if $Z \in \{c_1 H^1 + \dots + c_r H^r \mid c_1, \dots, c_r \in \{0, 1\}\}$ up to conjugation. This property is related to the structure of the attached nilmanifold.

Proposition 2.3 *Let $(\mathfrak{n}, B_\sigma)$ be the nilmanifold attached to a graded Lie algebra $\mathfrak{g} = \sum \mathfrak{g}_k$ of ν -th kind. Then,*

- (1) *there exists a type α_0 gradation $\mathfrak{g} = \sum \mathfrak{g}'_k$ which satisfies $\mathfrak{n} = \sum_{k>0} \mathfrak{g}'_k$,*
- (2) *$\mathfrak{n}$ is of ν -step nilpotent if and only if the gradation is of type α_0 .*

Proof. (1): Denote the characteristic element of the gradation $\mathfrak{g} = \sum \mathfrak{g}_k$ by $Z = c_{i_1}H^{i_1} + \dots + c_{i_s}H^{i_s}$ with $c_{i_1}, \dots, c_{i_s} > 0$. Then the gradation given by $Z' = H^{i_1} + \dots + H^{i_s}$ satisfies the condition. (2): Assume $\mathfrak{g} = \sum \mathfrak{g}_k$ is of type α_0 . Since $\mathfrak{g}_1$ generates $\mathfrak{g}_\nu$, the lower central series satisfies $C^\nu(\mathfrak{n}) \neq 0$. Thus $\mathfrak{n}$ is of ν -step nilpotent. To show the converse, assume $\mathfrak{g} = \sum \mathfrak{g}_k$ is not of type α_0 . In this case one has $Z = c_{i_1}H^{i_1} + \dots + c_{i_s}H^{i_s}$ and some coefficients are greater than 1. As in the proof for (1), the gradation $\mathfrak{g} = \sum \mathfrak{g}'_k$ given by $H^{i_1} + \dots + H^{i_s}$ is of type α_0 and satisfies $\mathfrak{n} = \sum_{k>0} \mathfrak{g}'_k$. The assumption implies that $\mathfrak{g} = \sum \mathfrak{g}'_k$ is of ν' -kind with $\nu' < \nu$. Therefore $\mathfrak{n}$ is ν' -step nilpotent and not ν -step nilpotent. $\square$

This proposition means that, to study the nilmanifold attached to GLA, it is enough to consider gradations of type α_0 .

3 Ricci curvatures of the attached nilmanifolds

In this section we calculate the Ricci curvatures of the nilmanifolds attached to GLA. The Ricci curvatures can be written by using the bracket products and the grade-reversing Cartan involutions.

Lemma 3.1 *The nilmanifolds $(\mathfrak{n}, B_\sigma)$ attached to GLA satisfy that*

- (1) $B_\sigma([U, V], W) = -B_\sigma(V, [\sigma U, W])$ for every $U, V, W \in \mathfrak{n}$,
- (2) $(\text{ad}_U)^*V = [V, \sigma U]_\mathfrak{n}$ for every $U, V \in \mathfrak{n}$.

These formulae can be checked easily but quite useful for calculating the Ricci curvatures.

Definition 3.2 Let $\{E_i^{(k)}\}$ be an orthonormal basis of $\mathfrak{g}_k$. We call $Z_k := \sum [\sigma(E_i^{(k)}), E_i^{(k)}]$ the k -th mean curvature vector.

It is easy to show that Z_k is independent of the choice of orthonormal basis. Furthermore, Lemma 3.1 leads that the vector $H_0 := Z_1 + \dots + Z_\nu$ coincides with the mean curvature vector of the solvmanifold $(\mathfrak{a} + \mathfrak{n}, B_\sigma)$.

Lemma 3.3 *Let $\{E_i^{(k)}\}$ be an orthonormal basis of $\mathfrak{g}_k$ and define the operators A_k and B_k on $\mathfrak{n}$ by*

$$A_k(U) := \sum [E_i^{(k)}, [\sigma E_i^{(k)}, U]], \quad B_k(U) := \sum [\sigma E_i^{(k)}, [E_i^{(k)}, U]].$$

Then, for every $U_l \in \mathfrak{g}_l$, we have

- (1) $A_k(U_l) = B_k(U_l) - [Z_k, U_l],$
- (2) $B_k(U_l) = A_{k+l}(U_l) = B_{k+l}(U_l) - [Z_{k+l}, U_l].$

Proof. (1) is a direct consequence of the Jacobi identity. To show (2), we need an orthonormal basis $\{E_j^{(k+l)}\}$ of $\mathfrak{g}_{k+l}$. It follows from $[E_i^{(k)}, U_l] \in \mathfrak{g}_{k+l}$ that

$$\begin{aligned} B_m(U_l) &= \sum_i [\sigma E_i^{(k)}, [E_i^{(k)}, U_l]] \\ &= \sum_i \sum_j [\sigma E_i, \langle [E_i^{(k)}, U_l], E_j^{(k+l)} \rangle E_j^{(k+l)}] \\ &= \sum_i \sum_j [\sigma (\langle E_i^{(k)}, [\sigma U_l, E_j^{(k+l)}] \rangle E_i^{(k)}), E_j^{(k+l)}] \\ &= \sum_j [\sigma ([\sigma U_l, E_j^{(k+l)}]), E_j^{(k+l)}] \\ &= A_{l+m}(U_l). \end{aligned}$$

This concludes the lemma. □

By definition, $B_k(U_l) = 0$ if $k + l > \nu$. Therefore one can determine A_k and B_k by these recursion formulae. These operators play important roles for calculating the Ricci curvatures of our nilmanifolds.

Theorem 3.4 *Let $(\mathfrak{n}, B_\sigma)$ be the nilmanifold attached to GLA and $\text{Ric}^{\mathfrak{n}}$ be the Ricci curvature. Then, for every $U_l \in \mathfrak{g}_l$, one has*

- (1) $\text{Ric}^{\mathfrak{n}}(U_l) = -(1/4) \sum_{m < l} A_m(U_l) + (1/2) \sum_{m > l} A_m(U_l),$
- (2) $\text{Ric}^{\mathfrak{n}}(U_1) = (1/2) \sum (1 - k) [Z_k, U_1],$
- (3) $\text{Ric}^{\mathfrak{n}}(U_2) = (1/4) \sum (2 - k) [Z_k, U_2].$

Proof. Let $\{E_i\}$ be an orthonormal basis of $\mathfrak{g}_k$. Direct calculations lead that

$$\sum \text{ad}_{E_i} \circ (\text{ad}_{E_i})^*(U_l) = - \sum [E_i, [\sigma E_i, U_l]_{\mathfrak{n}}] = \begin{cases} -A_m & \text{for } m < l, \\ 0 & \text{for } m \geq l. \end{cases}$$

Furthermore, one can see that

$$\sum (\text{ad}_{E_i})^* \circ \text{ad}_{E_i}(U_l) = -B_m(U_l) = -A_{l+m}(U_l).$$

Thus Proposition 1.4 concludes (1). One can calculate $A_k(U_1)$ and $A_k(U_2)$ by using Lemma 3.3 as follows:

$$\begin{aligned} A_k(U_1) &= B_k(U_1) - [Z_k, U_1] \\ &= (1/2)B_{k+1}(U_1) - [Z_k, U_1] - [Z_{k+1}, U_1] \\ &= \dots \\ &= -[Z_k + Z_{k+1} + \dots + Z_\nu, U_1], \\ A_k(U_2) &= -[Z_k, U_2] - [Z_{k+2}, U_2] - [Z_{k+4}, U_2] - \dots \end{aligned}$$

By substituting these formulae for the equation (1), we conclude (2) and (3). $\square$

4 Mean curvature vectors

As we saw in the previous section, one needs to know the mean curvature vectors $Z_1, \dots, Z_\nu$ for calculating the Ricci curvature of the nilmanifolds attached to GLA. In this section we investigate these vectors.

We call $H_\alpha \in \mathfrak{a}$ the *root vector* of a root α if $B_\sigma(H_\alpha, H) = \alpha(H)$ for every $H \in \mathfrak{a}$. Our root vectors are the negative of the usual ones, since $B_\sigma = -B$ on $\mathfrak{a} \times \mathfrak{a}$. We employ this definition for convenience. Let $\Delta_k := \{\alpha \in \Delta \mid \alpha(Z) = k\}$.

Lemma 4.1 *The k -th mean curvature vector Z_k satisfies (1) $Z_k = \sum_{\alpha \in \Delta_k} (\dim \mathfrak{g}_\alpha) H_\alpha$, and (2) $B_\sigma(Z_k, Z) = k \dim \mathfrak{g}_k$.*

Proof. Let $E \in \mathfrak{g}_\alpha$ be a unit vector. Then, Lemma 3.1 follows that $H_\alpha = [\sigma E, E]$, which concludes (1). To show (2) we need $\{E_i\}$, an orthonormal basis of $\mathfrak{g}_k$. One can see that

$$\begin{aligned} B_\sigma(Z_k, Z) &= B_\sigma\left(\sum [\sigma E_i, E_i], Z\right) = \sum B_\sigma(E_i, [Z, E_i]) \\ &= \sum B_\sigma(E_i, kE_i) = k \dim \mathfrak{g}_k. \end{aligned}$$

This concludes (2). $\square$

We now determine the mean curvature vectors of our nilmanifolds. Recall that we can assume, without loss of generality, that Z is a linear combination of $\{H^i\}$, the dual basis of the simple roots $\Pi = \{\alpha_i\}$. We also recall that the Weyl group is generated by $\{s_1, \dots, s_r\}$, where s_j denotes the reflection of $\mathfrak{a}$ with respect to the hyperplane $\{H \in \mathfrak{a} \mid \alpha_j(H) = 0\}$.

Theorem 4.2 *Let $(\mathfrak{n}, B_\sigma)$ be the nilmanifold attached to GLA and assume that the characteristic element is $Z = c_{i_1} H^{i_1} + c_{i_2} H^{i_2} + \dots + c_{i_s} H^{i_s}$ with $c_{i_1}, \dots, c_{i_s} > 0$. Then,*

- (1) $Z_k = z_{i_1}H^{i_1} + \cdots + z_{i_s}H^{i_s}$ with $z_{i_j} = \sum_{\beta \in \Delta_k} (\dim \mathfrak{g}_\beta) B_\sigma(H_{\alpha_{i_j}}, H_\beta)$, for every k , and
- (2) $H_0 = h_{i_1}H^{i_1} + \cdots + h_{i_s}H^{i_s}$ with $h_{i_j} = \sum_{\beta \in \Delta^{i_j}} (\dim \mathfrak{g}_\beta) B_\sigma(H_{\alpha_{i_j}}, H_\beta)$, where $\Delta^{i_j} := \{\beta \in \Delta \mid \beta(Z) = \beta(H^{i_j}) > 0\}$. Furthermore, $h_{i_j} > 0$.

Proof. (1): First of all we show that $Z_k \in V := \mathbb{R}H^{i_1} + \mathbb{R}H^{i_2} + \cdots + \mathbb{R}H^{i_s}$. Since the orthogonal complement of V in $\mathfrak{a}$ is spanned by root vectors $\{H_{\alpha_j} \mid j \neq i_1, \dots, i_s\}$, it is sufficient to show that $B_\sigma(Z_k, H_{\alpha_j}) = 0$ for every $j \neq i_1, \dots, i_s$. By the definition of Weyl group, this condition is equivalent to $s_j(Z_k) = Z_k$. Take $j \neq i_1, \dots, i_s$. Because s_j acts trivially on V , one has $s_j(Z) = Z$ and hence $s_j(\Delta_k) = \Delta_k$. This and Lemma 4.1 imply that

$$s_j(Z_k) = s_j\left(\sum_{\beta \in \Delta_k} (\dim \mathfrak{g}_\beta) H_\beta\right) = \sum_{\beta \in \Delta_k} (\dim \mathfrak{g}_\beta) s_j(H_\beta) = \sum_{\beta \in \Delta_k} (\dim \mathfrak{g}_{s_j(\beta)}) H_{s_j(\beta)} = Z_k.$$

Note that the multiplicities of roots, $\dim \mathfrak{g}_\beta$, are invariant under the action of the Weyl group. Therefore we have $B_\sigma(Z_k, H_{\alpha_j}) = 0$, and hence $Z_k \in V$. Each coefficient of $Z_k = z_{i_1}H^{i_1} + \cdots + z_{i_s}H^{i_s}$ can be determined by

$$z_{i_j} = \alpha_{i_j}(Z_k) = \alpha_{i_j}\left(\sum_{\beta \in \Delta_k} (\dim \mathfrak{g}_\beta) H_\beta\right) = \sum_{\beta \in \Delta_k} (\dim \mathfrak{g}_\beta) B_\sigma(H_{\alpha_{i_j}}, H_\beta).$$

(2): Let $\Delta' := \{\beta \in \Delta \mid \beta(Z) > 0\}$. Because of (1), one can write $H_0 = h_{i_1}H^{i_1} + \cdots + h_{i_s}H^{i_s}$ with $h_{i_j} = \sum_{\beta \in \Delta'} (\dim \mathfrak{g}_\beta) B_\sigma(H_{\alpha_{i_j}}, H_\beta)$. We here claim that s_{i_j} preserves $\Delta' - \Delta^{i_j}$. For every $\beta \in \Delta' - \Delta^{i_j}$, one can write $\beta = \sum n_k \alpha_k$ so that $n_{i_t} > 0$ for some $t \neq j$. The formula of the reflection

$$s_{i_j}(\beta) = \beta - 2 \frac{\langle \alpha_{i_j}, \beta \rangle}{\langle \beta, \beta \rangle} \alpha_{i_j}$$

leads that the α_{i_t} -coefficient of $s_{i_j}(\beta)$ is also n_{i_t} . Hence, $s_{i_j}(\beta)(Z) \geq n_{i_t} > 0$ and $s_{i_j}(\beta)(Z) \geq n_{i_t} + \beta(H^{i_j}) > \beta(H^{i_j})$. This completes the claim. Thus, the similar argument as (1) leads that

$$\sum_{\beta \in \Delta' - \Delta^{i_j}} (\dim \mathfrak{g}_\beta) B_\sigma(H_{\alpha_{i_j}}, H_\beta) = 0,$$

which concludes the first part of (2). Now we show that $h_{i_j} > 0$. There might exist $\beta \in \Delta^{i_j}$ such that $B_\sigma(H_{\alpha_{i_j}}, H_\beta) < 0$. But, for such β , one has $s_{i_j}(\beta) \in \Delta^{i_j}$ and $B_\sigma(H_{\alpha_{i_j}}, s_{i_j}(H_\beta)) = -B_\sigma(H_{\alpha_{i_j}}, H_\beta)$. Therefore every negative term cancels with another term. Hence, we got $h_{i_j} = \sum (\dim \mathfrak{g}_\beta) B_\sigma(H_{\alpha_{i_j}}, H_\beta)$, where β run through $\{\beta \in \Delta^{i_j} \mid B_\sigma(H_{\alpha_{i_j}}, H_\beta) > 0, s_{i_j}(\beta) \notin \Delta^{i_j}\}$. Since this set contains α_{i_j} , one has $h_{i_j} > 0$. $\square$

Next we show a remarkable property of mean curvature vectors, which is quite useful for calculating the Ricci curvatures of our nilmanifolds and solvmanifolds.

Theorem 4.3 *Let $(\mathfrak{n}, B_\sigma)$ be the nilmanifold attached to GLA with the characteristic element Z . Then the mean curvature vectors Z_k satisfy*

$$2 \sum_{k>0} k Z_k = Z.$$

Proof. It is enough to show that $\langle 2 \sum k Z_k, H \rangle = \langle Z, H \rangle$ for every $H \in \mathfrak{a}$. Let E_i be an orthonormal basis of $\mathfrak{g}$. By the definition of the Killing form, one has

$$\langle Z, H \rangle = \text{tr}(\text{ad}_Z \circ \text{ad}_H) = \sum_i \langle [Z, [H, E_i]], E_i \rangle.$$

One can assume that each E_i is contained in the root spaces. Denote by $E_\alpha \in \mathfrak{g}_\alpha$ a unit vector. Thus we have

$$\begin{aligned} \langle Z, H \rangle &= \sum_{\alpha \in \Delta} \langle [Z, [H, E_\alpha]], E_\alpha \rangle \dim \mathfrak{g}_\alpha \\ &= \sum_{\alpha \in \Delta} \alpha(Z) \alpha(H) \dim \mathfrak{g}_\alpha. \end{aligned}$$

On the other hand, Lemma 4.1 (1) leads that

$$2 \sum k Z_k = 2 \sum_{\alpha \in \Delta^+} \alpha(Z) (\dim \mathfrak{g}_\alpha) H_\alpha = \sum_{\alpha \in \Delta} \alpha(Z) (\dim \mathfrak{g}_\alpha) H_\alpha.$$

Therefore one can easily see that

$$\langle 2 \sum k Z_k, H \rangle = \sum_{\alpha} \alpha(Z) (\dim \mathfrak{g}_\alpha) \alpha(H),$$

which completes the proof. $\square$

Combining this theorem with Theorem 3.4, the Ricci curvatures of our nilmanifolds can be written easier.

Corollary 4.4 *For the nilmanifold $(\mathfrak{n}, B_\sigma)$ attached to GLA, we have*

- (1) $\text{Ric}^{\mathfrak{n}}(U_1) = -(1/4)U_1 + (1/2) \sum [Z_k, U_1]$ for $U_1 \in \mathfrak{g}_1$,
- (2) $\text{Ric}^{\mathfrak{n}}(U_2) = -(1/4)U_2 + (1/2) \sum [Z_k, U_2]$ for $U_2 \in \mathfrak{g}_2$.

5 Curvature properties of the solvable extensions

We now consider one-dimensional solvable extensions of the nilmanifolds $(\mathfrak{n}, B_\sigma)$ attached to GLA, and investigate their curvature properties and Einstein conditions. We consider the following solvable extensions:

$$\mathfrak{s} := \mathbb{R}H + \mathfrak{n}, \quad \langle, \rangle^c := cB_\sigma|_{\mathbb{R}H \times \mathbb{R}H} + B_\sigma|_{\mathfrak{n} \times \mathfrak{n}},$$

where $c > 0$, $H \in \mathfrak{a}$ and $\text{ad}_H|_{\mathfrak{n}}$ has only positive eigenvalues.

A solvmanifold $(\mathfrak{s}, \langle, \rangle)$ is called a *Carnot space* if $\dim[\mathfrak{s}, \mathfrak{s}]^\perp = 1$ and there exists $A_0 \in [\mathfrak{s}, \mathfrak{s}]^\perp$ such that every eigenvalue of ad_{A_0} on $[\mathfrak{s}, \mathfrak{s}]$ is a positive integer (see [19]). We call a Carnot space is of $(k+1)$ -step if k is the maximum of the eigenvalues of ad_{A_0} .

Proposition 5.1 *The above solvmanifold $(\mathfrak{s} = \mathbb{R}H + \mathfrak{n}, \langle, \rangle^c)$ is of Iwasawa-type. This solvmanifold is a Carnot space if every eigenvalue of $\text{ad}_{aH}|_{\mathfrak{n}}$ is a positive integer for some $a > 0$.*

The Ricci curvatures of solvable extensions can be calculated as follows.

Lemma 5.2 *Denote the Ricci curvatures of $(\mathfrak{s} = \mathbb{R}H + \mathfrak{n}, \langle, \rangle^c)$ and $(\mathfrak{n}, B_\sigma)$ by Ric^c and $\text{Ric}^{\mathfrak{n}}$, respectively. Let ad and $\text{ad}^{\mathfrak{g}}$ be the adjoint operators of $\mathfrak{s}$ and $\mathfrak{g}$, respectively. We normalize H to be the mean curvature vector of $(\mathfrak{s}, \langle, \rangle^1 = B_\sigma)$. Then,*

- (1) $\text{Ric}^c(H) = -(\text{tr}(\text{ad}_H)^2)/(c \cdot \text{tr}(\text{ad}_H^{\mathfrak{g}})^2) \cdot H$,
- (2) if $[H, \mathfrak{g}_0] = 0$, then $\text{Ric}^c(H) = -(1/2c)H$,
- (3) $\text{Ric}^c(U) = \text{Ric}^{\mathfrak{n}}(U) - (1/c)[H, U]$ for every $U \in \mathfrak{n}$,
- (4) $\text{Ric}^c(U_1) = -(1/4)U_1 + [(1/2) \sum Z_k - (1/c)H, U_1]$ for every $U_1 \in \mathfrak{g}_1$,
- (5) $\text{Ric}^c(U_2) = -(1/4)U_2 + [(1/2) \sum Z_k - (1/c)H, U_2]$ for every $U_2 \in \mathfrak{g}_2$.

Proof. (1): Proposition 1.3 leads that $\text{Ric}^c(H) = -(\text{tr}(\text{ad}_H)^2/|H|^2) \cdot H$. By the definition of the inner product, we have

$$|H|^2 = c \cdot B_\sigma(H, H) = c \cdot \text{tr}(\text{ad}_H^{\mathfrak{g}})^2.$$

(2): If $[H, \mathfrak{g}_0] = 0$, then we have

$$\text{tr}(\text{ad}_H^{\mathfrak{g}})^2 = \text{tr}(\text{ad}_H^{\mathfrak{g}}|_{\mathfrak{n}})^2 + \text{tr}(\text{ad}_H^{\mathfrak{g}}|_{\mathfrak{g}_0})^2 + \text{tr}(\text{ad}_H^{\mathfrak{g}}|_{\sigma(\mathfrak{n})})^2 = 2\text{tr}(\text{ad}_H)^2.$$

We finish the proof of (2) by substituting this for (1).

(3): Note that the mean curvature vector of $(\mathfrak{s}, \langle, \rangle^c)$ is $(1/c)H$. Thus Proposition 1.3 leads that

$$\text{Ric}^c(U) = \text{Ric}^{\mathfrak{n}}(U) - (\text{tr}(\text{ad}_{(1/c)H})/|(1/c)H|^2) \cdot [(1/c)H, U]$$

for $U \in \mathfrak{n}$. Recall that $\langle H, A \rangle^1 = \text{tr}(\text{ad}_A)$ for every $A \in \mathbb{R}H$, since H is the mean curvature vector of $(\mathfrak{s}, \langle, \rangle^1)$. Thus one has

$$|(1/c)H|^2 = \langle (1/c)H, (1/c)H \rangle^c = (1/c)\langle H, H \rangle^1 = (1/c)\text{tr}(\text{ad}_H) = \text{tr}(\text{ad}_{(1/c)H}).$$

This completes the proof of (3).

(4) and (5): They are direct consequences of (3) and Corollary 4.4. $\square$

Next we study the Einstein condition of the solvable extension. The following theorem implies that there is the only one natural choice of the solvable extension.

Theorem 5.3 *Let $(\mathfrak{n}, \langle, \rangle)$ be the nilmanifold attached to a graded Lie algebra of type α_0 and consider the above solvable extension $(\mathfrak{s} = \mathbb{R}H + \mathfrak{n}, \langle, \rangle^c)$. Then we have the following:*

- (1) *If $(\mathfrak{s}, \langle, \rangle^c)$ is Einstein, then $\text{Ric}^c(H) = -(1/2c)H$.*
- (2) *Assume that $\mathfrak{n}$ is abelian. Then $(\mathfrak{s}, \langle, \rangle^c)$ is Einstein if and only if $H \parallel H_0$.*
- (3) *Assume that $\mathfrak{n}$ is not abelian. If $(\mathfrak{s}, \langle, \rangle^c)$ is Einstein, then $c = 2$ and $H \parallel H_0$.*

Proof. First of all, we normalize H to be the mean curvature vector of $(\mathfrak{s}, \langle, \rangle^1 = B_\sigma)$.

(1): Assume that $(\mathfrak{s}, \langle, \rangle^c)$ is Einstein with Einstein constant s . We have only to show $[H, \mathfrak{g}_0] = 0$, because of Lemma 5.2 (2). Let $U_1 \in \mathfrak{g}_1$. Einstein condition and Lemma 5.2 (4) lead that

$$[-(1/4)Z + (1/2) \sum Z_k - (1/c)H, U_1] = \text{Ric}^c(U_1) = sU_1 = [sZ, U_1].$$

Thus, the element $U := -(1/4)Z + (1/2) \sum Z_k - (1/c)H - sZ$ acts trivially on $\mathfrak{g}_1$. By assumption, the gradation is of type α_0 , that is, $\mathfrak{g}$ is generated by $\mathfrak{g}_1$ and the grade-reversing Cartan involution. Therefore, $\sigma(U) = -U$ and $[U, \mathfrak{g}_1] = 0$ lead that $[U, \mathfrak{g}] = 0$. We conclude that

$$-(1/4)Z + (1/2) \sum Z_k - (1/c)H = sZ, \quad (5.1)$$

because $\mathfrak{g}$ is semisimple. Theorem 4.2 leads that $[Z, \mathfrak{g}_0] = 0$ and $[Z_k, \mathfrak{g}_0] = 0$, which conclude $[H, \mathfrak{g}_0] = 0$.

(2): We consider the case that $\mathfrak{n}$ is abelian. Note that, in this case, we have $H_0 = \sum Z_k = Z_1 = (1/2)Z$ (the last equality follows from Theorem 4.3). If $(\mathfrak{s}, \langle, \rangle^c)$ is Einstein with Einstein constant s , the equation (5.1) implies that H is parallel to H_0 . To show the converse, we assume that $H \parallel H_0$. One has $\text{Ric}^c(H_0) = -(1/2c)H_0$, because of $[H_0, \mathfrak{g}_0] = 0$ and Lemma 5.2 (2). One also has

$$\text{Ric}^c(U_1) = -(1/c)[H, U_1] = -(1/2c)[Z, U_1] = -(1/2c)U_1,$$

because of $\text{Ric}^n = 0$ and Lemma 5.2 (3). Therefore $(\mathfrak{s}, \langle, \rangle^c)$ is Einstein with Einstein constant $-(1/2c)$.

(3): We consider the case that $\mathfrak{n}$ is not abelian (that is, $\mathfrak{g}_2 \neq 0$) and $(\mathfrak{s}, \langle, \rangle^c)$ is Einstein. For $U_1 \in \mathfrak{g}_1$ and $U_2 \in \mathfrak{g}_2$, Einstein condition and Lemma 5.2 (4), (5) imply that

$$[(1/2) \sum Z_k - (1/c)H, U_1] = sU_1, \quad [(1/2) \sum Z_k - (1/c)H, U_2] = sU_2,$$

where the Einstein constant is $-(1/4) + s$. On the other hand, since our gradation is of type α_0 , one has $\mathfrak{g}_2 = [\mathfrak{g}_1, \mathfrak{g}_1]$. This implies that $(1/2) \sum Z_k - (1/c)H$ acts on $\mathfrak{g}_2$ as $2s \cdot \text{id}$. Therefore $s = 0$ and $H \parallel \sum Z_k = H_0$. Recall that we normalized H to be mean curvature vector, which leads that $H = H_0$ and $c = 2$. $\square$

In the case that $\mathfrak{n}$ is abelian, that is, the gradation is of first kind, the Einstein space $(\mathfrak{s}, \langle, \rangle^c)$ is obviously isometric to the real hyperbolic space, which has been well-known. In the case that $\mathfrak{n}$ is not abelian, our class of solvmanifolds contains many new examples of Einstein solvmanifolds. We will see examples in the latter sections.

6 Solvmanifolds attached to graded Lie algebras

Theorem 5.3 states that the following solvable extension is natural:

Definition 6.1 Let $(\mathfrak{n}, B_\sigma)$ be the nilmanifold attached to a graded Lie algebra and H_0 the mean curvature vector of $(\mathfrak{a} + \mathfrak{n}, B_\sigma)$. We call $(\mathfrak{s} = \mathbb{R}H_0 + \mathfrak{n}, \langle, \rangle^2)$ the *solvmanifold attached to a graded Lie algebra*.

Note that the solvmanifold attached to GLA is a Carnot space, because of Theorem 4.2. On the Ricci curvatures of these solvmanifolds, Lemma 5.2 immediately leads the following:

Theorem 6.2 *The Ricci curvature Ric of a solvmanifold attached to a graded Lie algebra, $(\mathfrak{s} = \mathbb{R}H_0 + \mathfrak{n}, \langle, \rangle^2)$, satisfies that*

- (1) $\text{Ric} = -(1/4)\text{id}$ on $\mathbb{R}H_0 + \mathfrak{g}_1 + \mathfrak{g}_2$,
- (2) $\text{Ric}(U) = \text{Ric}^{\mathfrak{n}}(U) - (1/2)[H_0, U]$ for every $U \in \mathfrak{n}$.

We here mention some relations between our solvmanifolds and symmetric spaces of noncompact type.

Let us consider $\mathfrak{g} = \sum_{k=-\nu}^{\nu} \mathfrak{g}_k$ the longest gradation. Here, we call the gradation *longest* if the characteristic element Z is a regular element. In this case, $\mathfrak{n} = \sum_{k>0} \mathfrak{g}_k = \sum_{\alpha \in \Delta^+} \mathfrak{g}_\alpha$ coincides with the nilpotent part of the Iwasawa decomposition of $\mathfrak{g}$. Furthermore, the solvmanifold $(\mathfrak{a} + \mathfrak{n}, \langle, \rangle)$, where $\langle, \rangle := 2B_\sigma|_{\mathfrak{a} \times \mathfrak{a}} + B_\sigma|_{\mathfrak{n} \times \mathfrak{n}}$, is nothing but the symmetric space of noncompact type, G/K , associated from $\mathfrak{g}$. This is a standard Einstein solvmanifold, whose rank one reduction coincides with our solvable extension $(\mathbb{R}H_0 + \mathfrak{n}, \langle, \rangle^2)$. Thus, the natural solvable extension is Einstein if the gradation is longest.

Let us consider the gradation $\mathfrak{g} = \sum_{k=-\nu}^{\nu} \mathfrak{g}_k$, which is not necessarily longest. In this case we have $\mathfrak{n} = \sum_{k>0} \mathfrak{g}_k \subset \sum_{\alpha \in \Delta^+} \mathfrak{g}_\alpha$. Therefore, our solvable Lie algebra $\mathbb{R}H_0 + \mathfrak{n}$ is a subalgebra of the solvable part of the Iwasawa decomposition. Thus one can see that our solvmanifold $(\mathbb{R}H_0 + \mathfrak{n}, \langle, \rangle^2)$ is a Riemannian submanifold in a symmetric space of noncompact type.

Because of the above observation, we have to investigate when our solvmanifold is totally geodesic, or the rank one reduction of a totally geodesic submanifold. As a result, this can only happen if the gradation is longest.

Proposition 6.3 *Let $\mathfrak{g} = \sum \mathfrak{g}_k$ be a gradation and $\mathfrak{n} := \sum_{k>0} \mathfrak{g}_k$. If $\mathfrak{n}$ is the nilradical of a totally geodesic submanifold, then the gradation is longest.*

Proof. Assume that $\mathfrak{a}' + \mathfrak{n}$ gives a totally geodesic submanifold and $\mathbb{R}H_0 + \mathfrak{n}$ is the rank one reduction of it. Denote by $\mathfrak{g} = \mathfrak{k} + \mathfrak{p}$ the Cartan decomposition and $\pi : \mathfrak{g} \rightarrow \mathfrak{p}$ the natural projection. One knows that $V := \pi(\mathfrak{a}' + \mathfrak{n})$ is a Lie triple system. If we denote by Z the characteristic element of our gradation, we have

$$V = \mathfrak{a}' + \sum_{\alpha(Z)>0} \mathfrak{p}_\alpha.$$

The totally geodesic submanifold coincides with the orbit of a Lie group with Lie algebra $[V, V] + V$.

We here claim that $[V, V] + V = \mathfrak{g}$. Let us take α such that $\alpha(Z) > 0$. Theorem 4.2 leads $\alpha(H_0) > 0$, and thus one has $[V, V] \supset [H_0, \mathfrak{p}_\alpha] = \mathfrak{k}_\alpha$. This implies $\mathfrak{g}_{-1} + \mathfrak{g}_1 \subset [V, V] + V$. Since we assumed that the gradation is of type α_0 , we conclude the claim.

Now it is easy to see from $\pi(\mathfrak{a}' + \mathfrak{n}) = \mathfrak{p}$ that the totally geodesic submanifold is the entire space, and Z is regular. $\square$

7 Two-step case

In this section we see that every solvmanifold attached to a graded Lie algebra of second kind is Einstein. Most of those Einstein solvmanifolds have the eigenvalue type $(1 < 2 ; d_1, d_2)$, but some of them have another eigenvalue type. We refer [13] for the explicit classification list of second kind graded Lie algebras.

Theorem 7.1 *Every solvmanifold $(\mathfrak{s}, \langle, \rangle)$ attached to a graded Lie algebra of second kind is Einstein.*

Proof. Since $\mathfrak{s} = \mathbb{R}H_0 + \mathfrak{g}_1 + \mathfrak{g}_2$, this is a direct consequence of Theorem 6.2. $\square$

We study their eigenvalue types. For a graded Lie algebra of second kind, $\mathfrak{g} = \sum_{k=-2}^2 \mathfrak{g}_k$, the characteristic element satisfies either $Z = H^i$ or $Z = H^i + H^j$. Recall that $\{H^i\}$ is the dual basis of the simple roots. If $Z = H^i$, Theorem 4.2 leads that $Z \parallel H_0$. In this case the eigenvalue type is $(1 < 2 ; d_1, d_2)$, by the definition of the characteristic element. If $Z = H^i + H^j$, Theorem 4.2 states only that $H_0 = aH^i + bH^j$. In this case unusual eigenvalue type may occur.

Let us consider the root system is of A_r -type. In this case $Z = H^i + H^j$ gives a second kind gradation. One can calculate the mean curvature vector directly,

$$H_0 = Z_1 + Z_2 = jH^i + (r + 1 - i)H^j,$$

up to scalar. If $i + j = r + 1$ then the eigenvalue type is $(1 < 2 ; d_1, d_2)$. If $i + j < r + 1$, the eigenvalue type is $(j < r + 1 - i < r + 1 - i + j ; d_1, d_2, d_3)$. These arguments imply that

Corollary 7.2 *For every $0 < \mu_1 < \mu_2$, there exists a standard Einstein solvmanifold $(\mathfrak{s}, \langle, \rangle)$ with two-step nilpotent $\mathfrak{n}$, whose eigenvalue type is $(\mu_1 < \mu_2 < \mu_1 + \mu_2 ; d_1, d_2, d_3)$.*

We see a simple example here. Consider the nilpotent subalgebra of $\mathfrak{sl}(n + m + l, \mathbb{R})$,

$$\mathfrak{n} := \left\{ \begin{pmatrix} 0 & X & Z \\ 0 & 0 & Y \\ 0 & 0 & 0 \end{pmatrix} \mid X \in M_{n,m}(\mathbb{R}), Y \in M_{m,l}(\mathbb{R}), Z \in M_{n,l}(\mathbb{R}) \right\}.$$

Let us consider the gradation of $\mathfrak{sl}(n + m + l, \mathbb{R})$ of second kind, given by the above block decomposition. Then $(\mathfrak{n}, B_\sigma)$ coincides with the nilmanifold attached to this gradation, therefore it has an Einstein solvable extension.

In this case, the root system is of $A_{n+m+l-1}$ -type, and direct calculations lead that

$$\begin{aligned} H^n &= \frac{1}{n + m + l} \begin{pmatrix} (m + l) \cdot 1_n & & \\ & -n \cdot 1_m & \\ & & -n \cdot 1_l \end{pmatrix}, \\ H^m &= \frac{1}{n + m + l} \begin{pmatrix} l \cdot 1_n & & \\ & l \cdot 1_m & \\ & & -(n + m) \cdot 1_l \end{pmatrix}. \end{aligned}$$

The characteristic element is $Z = H^n + H^m$. The solvmanifold $\mathbb{R}Z + \mathfrak{n}$ is a three-step Carnot space, but this is not what we want. Our Einstein solvable extension is $\mathbb{R}H_0 + \mathfrak{n}$, where H_0 is parallel to $(n + m)H^n + (m + l)H^m$. This is $(n + 2m + l + 1)$ -step Carnot space if $n + m$ and $m + l$ have no common divisor, whereas $\mathfrak{n}$ is two-step nilpotent.

We also note that, in case $n = l = 1$, this nilmanifold is nothing but the Heisenberg Lie algebra, and the constructed Einstein solvmanifold is isometric to the complex hyperbolic space. In case $n = l$, the eigenvalue type of the Einstein solvmanifold is $(1 < 2; nm + ml, nl)$. In case $n \neq l$, ad_{H_0} has three distinct eigenvalues.

8 Examples of Einstein spaces with higher steps

In this section we investigate the solvmanifolds attached to graded Lie algebras of third kind and of fourth kind.

Proposition 8.1 *Assume a graded Lie algebra is of third kind and each Z_k is parallel to Z . Then the attached solvmanifold $(\mathfrak{s}, \langle, \rangle)$ is Einstein if and only if $\dim \mathfrak{g}_1 = 2 \dim \mathfrak{g}_2$.*

Proof. Recall that Theorem 6.2 states $\text{Ric} = -(1/4) \cdot \text{id}$ on $\mathbb{R}H_0 + \mathfrak{g}_1 + \mathfrak{g}_2$. Therefore $(\mathfrak{s}, \langle, \rangle)$ is Einstein if and only if $\text{Ric}(U_3) = -(1/4)U_3$ for every $U_3 \in \mathfrak{g}_3$. One can calculate this as follows:

$$\begin{aligned}
\text{Ric}^n(U_3) &= (1/4)[Z_1, U_3] + (1/4)[Z_2, U_3], \\
\text{Ric}(U_3) &= \text{Ric}^n(U_3) - (1/2)[Z_1 + Z_2 + Z_3, U_3] \\
&= -(1/4)[Z_1, U_3] - (1/4)[Z_2, U_3] - (1/2)[Z_3, U_3] \\
&= -(1/4)((\dim \mathfrak{g}_1 + 2 \dim \mathfrak{g}_2 + 6 \dim \mathfrak{g}_3)/B_\sigma(Z, Z)) \cdot [Z, U_3] \\
&= -(1/4)(3 \dim \mathfrak{g}_1 + 6 \dim \mathfrak{g}_2 + 18 \dim \mathfrak{g}_3)/B_\sigma(Z, Z)) \cdot U_3.
\end{aligned}$$

Thus, $\text{Ric}(U_3) = -(1/4)U_3$ if and only if

$$3 \dim \mathfrak{g}_1 + 6 \dim \mathfrak{g}_2 + 18 \dim \mathfrak{g}_3 = B_\sigma(Z, Z) = 2(\dim \mathfrak{g}_1 + 4 \dim \mathfrak{g}_2 + 9 \dim \mathfrak{g}_3),$$

that is, $\dim \mathfrak{g}_1 = 2 \dim \mathfrak{g}_2$. □

Next we state the similar result for four-step case. Proof is also similar.

Proposition 8.2 *Assume a graded Lie algebra is of fourth kind and each Z_k is parallel to Z . Then the attached solvmanifold $(\mathfrak{s}, \langle, \rangle)$ is Einstein if and only if $\dim \mathfrak{g}_1 = 3 \dim \mathfrak{g}_3$ and $\dim \mathfrak{g}_1 + 4 \dim \mathfrak{g}_4 = 2 \dim \mathfrak{g}_2$.*

These propositions provide examples of Einstein solvmanifolds whose nilradicals are of three-step or four-step.

Theorem 8.3 *The solvmanifolds attached to the following graded Lie algebras are Einstein. The eigenvalue types of the constructed Einstein solvmanifolds are as follows.*

$\mathfrak{g}$	Δ	Z	eigenvalue type
$\mathfrak{g}_{2(2)}$	G_2	H^1	$(1 < 2 < 3 ; 2, 1, 2)$
$\mathfrak{g}_2^{\mathbb{C}}$	G_2	H^1	$(1 < 2 < 3 ; 4, 2, 4)$
$\mathfrak{f}_{4(4)}$	F_4	H^2	$(1 < 2 < 3 ; 12, 6, 2)$
$\mathfrak{f}_4^{\mathbb{C}}$	F_4	H^2	$(1 < 2 < 3 ; 24, 12, 4)$
$\mathfrak{e}_{6(2)}$	F_4	H^2	$(1 < 2 < 3 ; 18, 9, 2)$
$\mathfrak{e}_{7(-5)}$	F_4	H^2	$(1 < 2 < 3 ; 30, 15, 2)$
$\mathfrak{e}_{8(-24)}$	F_4	H^2	$(1 < 2 < 3 ; 54, 27, 2)$
$\mathfrak{e}_{6(6)}$	E_6	H^4	$(1 < 2 < 3 ; 18, 9, 2)$
$\mathfrak{e}_6^{\mathbb{C}}$	E_6	H^4	$(1 < 2 < 3 ; 36, 18, 4)$
$\mathfrak{f}_{4(4)}$	F_4	H^3	$(1 < 2 < 3 < 4 ; 6, 9, 2, 3)$
$\mathfrak{f}_4^{\mathbb{C}}$	F_4	H^3	$(1 < 2 < 3 < 4 ; 12, 18, 4, 6)$
$\mathfrak{e}_{6(2)}$	F_4	H^3	$(1 < 2 < 3 < 4 ; 12, 12, 4, 3)$
$\mathfrak{e}_{7(-5)}$	F_4	H^3	$(1 < 2 < 3 < 4 ; 24, 18, 8, 3)$
$\mathfrak{e}_{8(-24)}$	F_4	H^3	$(1 < 2 < 3 < 4 ; 48, 30, 16, 3)$

Proof. One has to check the relations of the dimensions. One can calculate $\dim \mathfrak{g}_k$ by counting roots and their multiplicities (one can see the table of multiplicities in [21]). $\square$

9 The case of low rank

In this section we mention the following problem: for a given semisimple Lie algebra $\mathfrak{g}$, is every attached solvmanifold Einstein? The answer is yes, if the rank of the associated symmetric space $(\mathfrak{g}, \mathfrak{k})$ is low.

If the rank of $(\mathfrak{g}, \mathfrak{k})$ is one, it is trivial. In this case, $\mathfrak{g}$ has only one gradation, which is longest, and the attached solvmanifold is the entire symmetric space.

Proposition 9.1 *Assume that the rank of $(\mathfrak{g}, \mathfrak{k})$ is two, that is, $\mathfrak{g}$ is one of the following:*

$$\mathfrak{sl}_3(\mathbb{R}), \mathfrak{sl}_3(\mathbb{C}), \mathfrak{su}^*(6), \mathfrak{e}_{6(-26)}, \mathfrak{so}(5, \mathbb{C}), \mathfrak{so}(2, n+2), \mathfrak{g}_{2(2)}, \mathfrak{g}_2^{\mathbb{C}}.$$

Then, for every gradation of $\mathfrak{g}$, the attached solvmanifold is Einstein.

Proof. Let $\{\alpha_1, \alpha_2\}$ be the set of simple roots. There are three gradations, given by H^1 , H^2 and $H^1 + H^2$. If the characteristic element is $H^1 + H^2$, the gradation is longest and thus the attached solvmanifold is Einstein. If the characteristic element is H^1 or H^2 , the gradation is of first kind, second kind or third kind. The solvmanifold attached to first kind or second kind gradation is Einstein (Theorems 6.2, 7.1). The third kind gradation is obtained by G_2 -type root system, whose attached solvmanifolds are Einstein by Theorem 8.3. $\square$

Similarly, we have the following:

Proposition 9.2 *If the root system of $(\mathfrak{g}, \mathfrak{k})$ is of A_3 -type, that is, $\mathfrak{g} = \mathfrak{sl}_4(\mathbb{R}), \mathfrak{sl}_4(\mathbb{C})$ or $\mathfrak{su}^*(8)$, every attached solvmanifold is Einstein.*

10 Rank one reduction of noncompact symmetric spaces

In this section we study the mean curvature vectors of noncompact symmetric spaces. For the graded Lie algebra with characteristic element $Z := H^1 + \cdots + H^r$, one knows that $\mathfrak{n}$ is the nilpotent part of the Iwasawa decomposition and the solvmanifold $(\mathfrak{a} + \mathfrak{n}, \langle, \rangle)$ is a symmetric space.

Proposition 10.1 *The mean curvature vector of the above solvmanifold $(\mathfrak{a} + \mathfrak{n}, \langle, \rangle)$ is $H_0 = h_1 H^1 + \cdots + h_r H^r$, where $h_i = (\dim \mathfrak{g}_{\alpha_i} + 4 \dim \mathfrak{g}_{2\alpha_i}) B_{\sigma}(H_{\alpha_i}, H_{\alpha_i})$.*

Proof. We apply Theorem 4.2 (2). In this case $\Delta^i := \{\beta \mid \beta(Z) = \beta(H^i) > 0\}$ consists of α_i and $2\alpha_i$ (if exists). Therefore,

$$h_i = (\dim \mathfrak{g}_{\alpha_i})B_\sigma(H_{\alpha_i}, H_{\alpha_i}) + (\dim \mathfrak{g}_{2\alpha_i})B_\sigma(H_{2\alpha_i}, H_{2\alpha_i}).$$

Proposition is completed by $H_{2\alpha_i} = 2H_{\alpha_i}$. □

This Proposition implies that, if every root has the same length, then H_0 is parallel to Z . It is remarkable that H_0 may not parallel to Z even for symmetric spaces. Furthermore, as we saw in Propositions 8.1 and 8.2, the Einstein condition requires some equalities of dimensions. It seem to be mysterious that symmetric spaces automatically satisfy these conditions.

References

- [1] D. V. Alekseevskii, Homogeneous Riemannian spaces of negative curvature, *Mat. Sb.* **25** (1975), 87-109; English translation, *Math. USSR-Sb.* **96** (1975), 93-117.
- [2] D. V. Alekseevskii, Classification of quaternionic spaces with transitive solvable group of motions, *Math. USSR-Izv.* **9** (1975), 297-339.
- [3] J. Berndt, F. Tricerri, L. Vanhecke, *Generalized Heisenberg groups and Damek-Ricci harmonic spaces*, Lect. Notes in Math. **1598** (1995), Springer-Verlag Berlin Heidelberg.
- [4] A. Besse, *Einstein manifolds*, Ergeb. Math. **10** (1987), Springer-Verlag, Berlin-Heidelberg.
- [5] J. Boggino, Generalized Heisenberg groups and solvmanifolds naturally associated, *Rend. Sem. Mat. Univ. Politec. Torino* **43** (1985), 529-547.
- [6] N. Bourbaki, *Groupes et algèbres de Lie, Chapitre IV - VI*, Éléments de Mathématique, Masson, Paris, 1981.
- [7] V. Cortés, Alekseevskian spaces, *Diff. Geom. Appl.* **6** (1996), 129-168.
- [8] E. Damek, F. Ricci, Harmonic analysis on solvable extensions of H -type groups. *J. Geom. Anal.* **2** (1992), 213-248.
- [9] P. Eberlien, J. Heber, Quarter pinched homogeneous spaces of negative curvature, *Internat. J. Math.* **7** (1996), 441-500.
- [10] C. S. Gordon, M. Kerr, New homogeneous Einstein metrics of negative Ricci curvature, *Ann. Global Anal. Geom.* **19** (2001), 75-101.

- [11] J. Heber, Noncompact homogeneous Einstein spaces, *Invent. Math.* **133** (1998), 279–352.
- [12] S. Helgason, *Differential geometry, Lie groups, and symmetric spaces*, Graduate Studies in Mathematics **34**, American Mathematical Society, Providence, RI, 2001.
- [13] S. Kaneyuki, On the subalgebras $\mathfrak{g}_0$ and $\mathfrak{g}_{\text{ev}}$ of semisimple graded Lie algebras, *J. Math. Soc. Japan* **45** (1993), 1–19.
- [14] S. Kaneyuki, H. Asano, Graded Lie algebras and generalized Jordan triple systems, *Nagoya Math. J.* **112** (1988), 81–115.
- [15] J. Lauret, Ricci soliton homogeneous nilmanifolds, *Math. Ann.* **319** (2001), 715–733.
- [16] J. Lauret, Finding Einstein solvmanifolds by a variational method, *Math. Z.* **241** (2002), 83–99.
- [17] J. Lauret, Minimal metrics on nilmanifolds, *Diff. Geom. Appl. Proc. Conf. Prague*, August 30 – September 3, 2004. Charles University, Prague (Czech Republic), 2005, 79–97.
- [18] K. Mori, Einstein metrics on Boggino-Damek-Ricci-type solvable Lie groups, *Osaka J. Math.* **39** (2002), 345–362.
- [19] S. Nishikawa, Harmonic maps and negatively curved homogeneous spaces, *Geometry and topology of submanifolds X* (Eds. W. H. Chen et. al.), World Sci. Publishing, Singapore, 2000, 200–215.
- [20] I. I. Pyateskii-Shapiro, *Automorphic functions and the geometry of classical domains*, Gordon and Breach Science Publishers, New York-London-Paris, 1969.
- [21] H. Tamaru, The local orbit types of symmetric spaces under the actions of the isotropy subgroups, *Diff. Geom. Appl.* **11** (1999), 29–38.
- [22] H. Tamaru, On certain subalgebras of graded Lie algebras, *Yokohama Math. J.* **46** (1999), 127–138.
- [23] H. Tamaru, A class of noncompact homogeneous Einstein manifolds, *Diff. Geom. Appl. Proc. Conf. Prague*, August 30 – September 3, 2004. Charles University, Prague (Czech Republic), 2005, 119–127.
- [24] T. H. Wolter, Einstein metrics on solvable Lie groups, *Math. Z.* **206** (1991), 457–471.